Applicants

Maureen J. Charron and Ellen B. Katz

Serial No. Filed 09/886,954 June 21, 2001

Page 3

Amendments to the Claims:

Please amend claims 1 and 11 as indicated below.

- 1. (Currently amended) A method for determining whether a subject has a defect in cell proliferation, comprising assaying a diagnostic sample of the subject for GLUTx expression, wherein detection of GLUTx expression elevated above normal is diagnostic of a defect in cell proliferation and wherein GLUTx protein het-subject for glutter as a subject has a diagnostic sample of the subject for GLUTx expression, wherein detection of GLUTx expression elevated above normal is diagnostic of a defect in cell proliferation and wherein GLUTx protein het-subject for glutter for g
- 2. (Original) The method of Claim 1, wherein the defect in cell proliferation is a neoplasm or a pre-neoplastic lesion.
- 3. (Original) The method of Claim 2, wherein the neoplasm is an adenocarcinoma.
- 4. (Original) The method of Claim 1, wherein the diagnostic sample is assayed using an agent reactive with GLUTx.
- 5. (Original) The method of Claim 4, wherein the agent is labeled with a detectable marker.
 - 6. (Original) The method of Claim 4, wherein the agent is an antibody.
- 7. (Original) The method of Claim 6, wherein the antibody is labeled with a detectable marker.



Applicants

Maureen J. Charron and Ellen B. Katz

Serial No.

09/886,954

Filed

June 21, 2001

Page 4

- 8. (Original) The method of Claim 1, wherein the diagnostic sample is assayed using at least one nucleic acid probe which hybridizes to nucleic acid encoding GLUTx.
- 9. (Original) The method of Claim 8, wherein the nucleic acid probe is DNA or RNA.
- 10. (Original) The method of Claim 9, wherein the nucleic acid probe is labeled with a detectable marker.

B1

- defect in cell proliferation in a subject who has undergone or is undergoing treatment for a defect in cell proliferation, comprising assaying a diagnostic sample of the subject for GLUTx expression, wherein detection of GLUTx expression elevated above normal in the diagnostic sample is indicative of a need to continue therapy to treat the defect in cell proliferation, and normal GLUTx expression in the diagnostic sample is indicative of successful therapy, and wherein GLUTx protein has comprises the amino acid sequence set forth in SEO ID NO:1 Genbank accession number XP_011828.
- 12. (Original) The method of Claim 11, wherein the defect in cell proliferation is a neoplasm or a pre-neoplastic lesion.
- 13. (Original) The method of Claim 12, wherein the neoplasm is an adenocarcinoma.
- 14. (Original) The method of Claim 11, wherein the diagnostic sample is assayed using an agent reactive with GLUTx.

١,

Applicants

Maureen J. Charron and Ellen B. Katz

Serial No. Filed 09/886,954 June 21, 2001

Page 5

15. (Original) The method of Claim 14, wherein the agent is labeled with a detectable marker.

- 16. (Original) The method of Claim 14, wherein the agent is an antibody.
- 17. (Original) The method of Claim 16, wherein the antibody is labeled with a detectable marker.
- 18. (Original) The method of Claim 11, wherein the diagnostic sample is assayed using at least one nucleic acid probe which hybridizes to nucleic acid encoding GLUTx.
- 19. (Original) The method of Claim 18, wherein the nucleic acid probe is DNA or RNA.
- 20. (Original) The method of Claim 19, wherein the nucleic acid probe is labeled with a detectable marker.
 - 21.-35. (Canceled)